

Deep Convective Cloud Calibration

D. R. Doelling, A. V. Gambheer

Analytical Services and Materials, Inc., Hampton, VA 23666

J. C. Stassi

SAIC, Hampton, VA 23666

L. Nguyen, D. F. Young, Y. Hu, P. Minnis,


NASA Langley Research Center, Hampton, VA

Matched angle technique

- Transfer calibration of MODIS (calibrated in the visible by solar diffuser) to Geostationary satellite (uncalibrated) by matching coincident, co-located, co-angled radiances
 - Bin visible pixel level radiances into 0.5° latitude by longitude regions (for GGO 1° regions)
 - Time match within 15 minutes
 - Angle match within 5° view angle and 15° azimuth (for GGO 10° view and 45° azimuth)


GOES-8 vs Terra-MODIS

monthly regression for Jan 2003






Force fit relies
on a fixed offset
and the mean

GOES-8 vs Terra-MODIS timeline



3 way matched angle cross-calibration validation



VIRS
GOES-8
0.9730

MODIS
VIRS
1.012

=


MODIS
GOES-8
.9847

From
timeline
0.9887

difference
0.4%

Jan 2001


GGEO noon cross-calibration regions



GOES-10 to GOES-8 noon cross-calibration during 2000-2003


GOES-10 vs GOES-8

Nov2000 (0.65um)






GOES-10 vs GOES-8

VIS, 0.65um




What happens operationally

- Matched angle conditions between 3 hourly geostationary and polar orbiter data tend to have preferred geographical locations and solar zenith angles that vary seasonally
- Navigation problems




Matched angle spatial sampling


Winter



Summer



GMS-5 navigational problems




Deep Convective Cloud Calibration

- Based on the assumption that there is maximum TOA albedo a cloud can have no matter how optically thick the cloud
- Can be used to detect relative gain drift in the visible sensor

Deep Convective Cloud Calibration



- At what optical depth does the TOA albedo stabilize, where cloud reflectivity and absorption balance?
- Fu-Liou radiative transfer code
 - McClatchey Tropical Profile
 - 10km thick ice cloud over ocean
 - 60 μ m particle size
 - Vary cloud optical depth from 1 to 1000



Deep convective
optical depths

Deep convective spectral albedo


Deep convective directional models



Deep convective cloud bidirectional models

- Hu Model
 - MODTRAN absorption
 - DISORT scattering
 - Ice cloud at 120 optical depth
 - Broadband model
- Terra Model
 - Bin CERES footprints into angular bins where the
 - MODIS 11 μ m Temperature < 205° K
 - Optical depth > 20
 - Ice cloud amount > 99%
- How lambertian are deep convective clouds?

Deep convective cloud bidirectional models





Test deep convective technique on VIRS data

- VIRS has a visible solar diffuser - should see no temporal drift if done correctly
- Use all pixels $< 205^{\circ}$ K
- Limit bidirectional effects
 - Limit solar zenith angle $< 40^{\circ}$ and view angle $< 40^{\circ}$
 - Bin visible radiances into 10° solar zenith, 10° view and 20° azimuth angle bins
 - Take mean of all sampled bins for a single precessionary (23 day) cycle (reduces the effects of preferential seasonally dependent angular bins)
- Use 4 precessionary cycles per year from 1998-2003
- Compare timeline from Hu and Terra model


VIRS deep convection timeline

HU

Terra





VIRS Visible radiance PDF for pixels with Temperature < 205°K




Sharpen PDF by using only pixels where the surrounding 8 pixels have a visible sigma < 8

- $\text{Wm}^{-2}\text{Sr}^{-1}\mu\text{m}^{-1}$ and a Temperature sigma < 1°K to reduce 3-D effects




Comparison of VIRS deep convective timeline techniques


Low sigma




Mode



Low sigma & Mode




VIRS deep convective dependencies



GGEO deep convective timelines using angular bin technique


GMS-5

COLD CLOUD GAIN for GMS-5




GOES-10

COLD CLOUD GAIN for GOES-10





GOES-8

COLD CLOUD GAIN for GOES-8




Comparison of GOES-8 calibrations

MODIS



VIRS



Conclusions

- Deep convective clouds can provide relative gain for non-calibrated visible instruments
 - Does not rely navigation
 - Does not need continuous monitoring of a calibrated visible source
 - Is well suited for operational calibration updating
- Use deep convective GGEO calibration as a sanity check of the current calibrations derived from MODIS in conjunction with noon GGEO cross-calibration